

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 June 2003 (05.06.2003)

PCT

(10) International Publication Number
WO 03/045332 A2

(51) International Patent Classification⁷: **A61K**

(21) International Application Number: PCT/US02/38258

(22) International Filing Date:
27 November 2002 (27.11.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/333,836 28 November 2001 (28.11.2001) US

(71) Applicant: **THE GENERAL HOSPITAL CORPORATION** [US/US]; 55 Fruit Street, Boston, MA 02114 (US).

(72) Inventor: **PODOLSKY, Daniel, K.**; 157 Edmunds Road, Wellesley, MA 02481 (US).

(74) Agents: **CLARK, Paul, T.** et al.; Clark & Elbing LLP, 101 Federal Street, Boston, MA 02110 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS AND COMPOSITIONS FOR TREATING LESIONS OF THE RESPIRATORY EPITHELIUM

(57) Abstract: This invention features methods of treating lesions of the airway epithelium by local or systemic administration of intestinal trefoil peptides. The intestinal trefoil peptide can be administered either alone or in combination with one or more therapeutic agents.

WO 03/045332 A2

METHODS AND COMPOSITIONS FOR TREATING LESIONS OF THE RESPIRATORY EPITHELIUM

Field of Invention

This invention relates to methods and compositions for treating lesions of the airway epithelium that can result, for example, from viral, bacterial, and fungal infections, inflammation, allergens, inhaled organic solvents, particulates, or irritant gases.

Background of the Invention

Upper airway lesions, including lesions from the external nasal nares to the larynx, are caused by a wide variety of local irritants, allergens, and infectious agents. Typically, these irritants give rise to the symptoms of rhinitis or 'runny nose.' In cases of severe lesions however, the tight junctions of the respiratory epithelial mucosa are disrupted such that entry of allergens or infectious agents is facilitated.

Tracheo-bronchial lesions (trachea and conducting bronchial tubes to the level of the respiratory bronchioles) are also commonly caused by respiratory infections, irritants, and allergens. Once the tracheo-bronchial epithelium and tight junctions have been disrupted, infectious, irritant, or allergic material may sensitize the lung, triggering the release of mediators, and subsequent airway constriction and asthma.

The alveolar epithelium, distal to the respiratory bronchioles, is generally well protected against infectious, irritant, and allergic exposure. However, infectious, immunologic, or chemical agents that penetrate the deep lung structures can cause pneumonias. Infectious agents that gain access to the systemic circulation in the lower airway can further result in sepsis pneumonias or a respiratory distress syndrome. Moreover, in certain inflammatory conditions

such as asthma, mucosal disruption results in increased levels of allergens and irritants, such that both inflammation and mucosal lysis are further exacerbated.

Rapid restoration of the normal airway epithelial barrier is therefore critical to reduce the damage caused by ongoing pathogenic or allergenic mechanisms in
5 respiratory tissues and alleviate the associated symptoms.

Summary of the Invention

The present invention features methods and compositions for the treatment of lesions of the airway epithelium in mammals, by administering to the mammal
10 therapeutically effective amounts of trefoil peptides, or a biologically active fragments thereof. Treatment of lesions according to the invention can speed healing, reduce pain, delay or prevent the occurrence of the lesion, and inhibit expansion, secondary infection, or other complications of the lesion. Lesions of the airway epithelium may result from any cause, including for example, an
15 allergic reaction, asthma, an infection, an inhaled chemical or particulate exposure, a thermal lesion, smoke inhalation, drug-induced lung damage, trauma (caused, for example, by surgery or intubation), a microbial infection (e.g., bacterial, viral, or fungal), chronic obstructive pulmonary disease, anti-neoplastic therapy, cystic fibrosis, cardiovascular compromise such as congestive heart
20 failure, or hyperbaric oxygen therapy.

In another aspect, the invention provides a composition, which includes a trefoil peptide in a pharmaceutically acceptable carrier suitable for inhalation administration. When formulated as such, the composition may be an aerosol (e.g., nasal spray, inhalation spray, inhalation solution, inhalation suspension)
25 administered by a metered dose inhaler. If desired, the formulation containing the trefoil peptide may be nebulized (e.g., by jet, ultrasonic nebulizer, or electronic nebulizer). Alternatively, the trefoil peptide formulation may be administered as a dry powder using a metered dose inhaler or a dry powder inhaler, for example.

In all foregoing aspects of the invention, the mammal is preferably a human and the trefoil peptide is human intestinal trefoil factor (ITF), spasmolytic peptide (SP), pS2, or biologically active fragments thereof. Such fragments include for example, ITF₁₅₋₇₃, ITF₂₁₋₇₃, ITF₁₋₇₂, ITF₁₅₋₇₂, or ITF₂₁₋₇₂ of SEQ ID NO.: 1

In the methods and compositions of this invention, a second therapeutic agent can be included. Such agents include anti-inflammatory agents such as glucocorticoids (beclomethasone, flunisolide, budesonide, triamcinolone, prednisolone, dexamethasone, or fluticasone) or non-steroidal anti-inflammatory agents (e.g., ibuprofen, tacrolimus, cromolyn, nedocromil, refecoxib, or celecoxib); antimicrobial agents (e.g., amikacin, gentamicin, kanamycin, neomycin, netilmicin, paromomycin, streptomycin, or tobramycin); antihistamines (e.g., diphenhydramine, fexofenadine, cetirizine, or loratadine); cholinergic receptor antagonists (e.g., ipratropium bromide or tiotropium); neurokinin receptor antagonists; leukotriene receptor antagonists; decongestants; phosphodiesterase inhibitors; or beta-adrenergic receptor antagonists (albuterol, bitolterol, epinephrine, fenoterol, formoterol, isoetharine, isoproterenol, metaproterenol, pirbuterol, procaterol, racepinephrine, salmeterol, or terbutaline). The second therapeutic agent may be administered within (either before or after) 14 days, 7 days, 1 day, 12 hours, 1 hour, or simultaneously with the trefoil peptide.

The second therapeutic agent can be present in the same or different pharmaceutical composition as the trefoil peptide. When the second therapeutic agent is present in a different pharmaceutical composition, different routes of administration may be used. For example, the second therapeutic agent may be administered orally, or by intravenous, intramuscular, or subcutaneous injection. Thus, the second therapeutic agent need not be administered by inhalation.

Of course, pharmaceutical compositions may contain two, three, or more trefoil peptides or biologically active trefoil peptide fragments. Alternatively, inhalation of the trefoil peptide may be supplemented by systemic (e.g., oral or

injectable) administration of the same or different trefoil peptide.

Airway epithelial lesions are prevented or ameliorated by administering the intestinal trefoil peptide-containing composition prior to the anticipated insult (e.g., surgery, or antineoplastic therapy for example). Preferably, the prophylactic
5 treatment begins at least one day, three days, five days, seven days, or ten days prior to the insult. Treatment of unanticipated airway lesions preferably begin immediately after insult, or within 24 hours.

In a preferred embodiment, the trefoil peptide or biologically active fragment is encoded by an isolated nucleic acid sequence that hybridizes under
10 high stringency conditions to a polynucleotide sequence having the sequence of SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, or SEQ ID NO: 9.

In another aspect, the invention features a pharmaceutical composition suitable for inhalation administration, containing a trefoil peptide or biologically
15 active fragment that is encoded by an isolated nucleic acid sequence that hybridizes under high stringency conditions to a polynucleotide sequence having the sequence of SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, or SEQ ID NO: 9

Mammalian trefoil peptides were discovered in 1982. One of the
20 mammalian trefoil peptides, human intestinal trefoil factor (ITF), has been characterized extensively, and is described in U.S. Patent Nos. 6,063,755, and 6,221,840, hereby incorporated by reference. The other two known human trefoil peptides are spasmolytic polypeptide (SP) and pS2. Trefoil peptides, described extensively in the literature (e.g., Sands *et al.*, Annu. Rev. Physiol. 58: 253-273
25 (1996), hereby incorporated by reference), are expressed in the gastrointestinal tract and have a three-loop structure formed by intrachain disulfide bonds between conserved cysteine residues. These peptides protect the intestinal tract from injury and can be used to treat intestinal tract disorders, such as peptic ulcers and inflammatory bowel disease. Homologs of these human peptides have been
30 found in a number of non-human animal species. All members of this protein

family, both human and non-human, are referred to herein as trefoil peptides.

Human ITF will be referred to most extensively in this application; however, the activity of human ITF is common to each of the mammalian trefoil peptides.

The term "trefoil peptide" is meant to include all mammalian homologs of human spasmolytic polypeptide, human pS2 and human ITF polypeptides, and biologically active fragments thereof. Homologs of the trefoil peptides have, preferably, 70% amino acid identity to the human sequence, more preferably 85% identity, most preferably 95%, or even 99% sequence identity. The length of comparison sequences will generally be at least about 8 amino acid residues, usually at least 20 amino acid residues, more usually at least 24 amino acid residues, typically at east 28 amino acid residues, and preferably more than 35 amino acid residues.

The term "fragment" is meant to include polypeptides that are truncations or deletions of SP, pS2 and ITF. Preferably, the fragments have 70% amino acid identity to the corresponding regions of the human polypeptide sequence. More preferably, the fragments are 85% identical, most preferably 95%, or even 99% identical to the human polypeptide sequence to which they correspond. The length of comparison sequences will generally be at least about 8 amino acid residues, usually at least 20 amino acid residues, more usually at least 24 amino acid residues, typically at east 28 amino acid residues, and preferably more than 35 amino acid residues.

Preferable fragments contain four cysteine residues in any positions which correspond to the cysteines at positions 25, 35, 45, 50, 51, 62, or 71, of human ITF (Figure 1), or positions 31, 41, 51, 56, 57, 68, and 82 of human pS2 (Figure 2). More preferably, fragments contain five cysteine residues at these positions. Most preferably, six, or even all seven cysteines are present.

Fragments of SP are meant to include truncations or deletions and preferably have 70% sequence identity to the corresponding human SP polypeptide sequence (Figure 3). More preferably, the fragments are 85% identical, most preferably 95%, or even 99% identical to the human polypeptide

sequence. Preferably, active fragments contain at least four cysteine residues, which correspond to positions 6, 8, 19, 29, 34, 35, 46, 58, 68, 78, 83, 84, 95, and 104 in the human SP polypeptide. More preferably, fragments contain six cysteines, which correspond to these positions. Even more preferable are
5 fragments that contain eight cysteines. Most preferable are fragments that contain cysteines at ten, twelve, or even, all fourteen positions.

It is recognized in the art that one function of the identified cysteine residues is to impart the characteristic three-loop (trefoil) structure. Accordingly, preferred fragments of ITF and pS2 have a least one loop structure, more
10 preferably, the fragments have two loop structures, and most preferably, they have three loop structures. It is equally well recognized that the native SP polypeptide has a six loop confirmation. Preferable fragments contain at least two of these loop structures, more preferably, four loop structures are conserved, and most preferably, five, or even all six loop structures are present.

15 By "aerosol" is meant any composition of the trefoil peptide of the invention administered as an aerosolized formulation, including for example an inhalation spray, inhalation solution, inhalation suspension, a nebulized solution, or nasal spray.

By "antimicrobial agent" is meant any compound that alters the growth of
20 bacteria or fungi cells, or viruses whereby growth is prevented, stabilized, or inhibited, or wherein the microbes are killed. In other words, the antimicrobial agents can be microbiocidal or microbiostatic.

By "antineoplastic therapy" is meant any treatment regimen used to treat cancer. Typical antineoplastic therapies include chemotherapy and radiation
25 therapy.

By "biologically active," when referring to a trefoil peptide, fragment, or homolog is meant any polypeptide that exhibits an activity common to its related, naturally occurring family member, and that the activity is common to the family of naturally occurring trefoil peptides. An example of a biological activity
30 common to the family of trefoil peptides is the ability to reconstitute the

gastrointestinal mucosa (Taupin *et al.*, Proc. Natl. Acad. Sci. U S A. 97(2): 799-804).

The term "isolated DNA" is meant DNA that is free of the genes which, in the naturally-occurring genome of the organism from which the given DNA is derived, flank the DNA. Thus, the term "isolated DNA" encompasses, for example, cDNA, cloned genomic DNA, and synthetic DNA.

The term "pharmaceutical composition" is meant any composition, which contains at least one therapeutically or biologically active agent and is suitable for administration to the patient. Pharmaceutical compositions suitable for delivering a therapeutic to the respiratory airways include, but are not limited to, aerosols and dry powders. Any of these formulations can be prepared by well-known and accepted methods of the art. See, for example, Remington: The Science and Practice of Pharmacy, 20th edition, (ed. AR Gennaro), Mack Publishing Co., Easton, PA, 2000.

By "high stringency conditions" is meant any set of conditions that are characterized by high temperature and low ionic strength and allow hybridization comparable with those resulting from the use of a DNA probe of at least 40 nucleotides in length, in a buffer containing 0.5 M NaHPO₄, pH 7.2, 7% SDS, 1mM EDTA, and 1% BSA (Fraction V), at a temperature of 65 C, or a buffer containing 48% formamide, 4.8X SSC, 0.2 M Tris-Cl, pH 7.6, 1X Denhardt's solution, 10% dextran sulfate, and 0.1% SDS, at a temperature of 42°C. Other conditions for high stringency hybridization, such as for PCR, Northern, Southern, or in situ hybridization, DNA sequencing, etc., are well known by those skilled in the art of molecular biology. See, e.g., F. Ausubel et al., Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY, 1998, hereby incorporated by reference. Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

By "substantially identical" is meant a polypeptide or nucleic acid exhibiting at least 75%, but preferably 85%, more preferably 90%, most preferably 95%, or 99% identity to a reference amino acid or nucleic acid

sequence. For polypeptides, the length of comparison sequences will generally be at least 20 amino acids, preferably at least 30 amino acids, more preferably at least 40 amino acids, and most preferably 50 amino acids. For nucleic acids, the length of comparison sequences will generally be at least 60 nucleotides, preferably at least 90 nucleotides, and more preferably at least 120 nucleotides.

By "therapeutically effective amount" is meant an amount sufficient to provide medical benefit. When administering trefoil peptides to a human patient according to the methods described herein, an effective amount will vary with the size of the lesion area being treated; however, a therapeutically effective amount is usually about 1-2500 mg of trefoil peptide per dose. Dosing is typically performed one to four times each day. The patient may also be administered with a trefoil peptide continuously over a set period of time.

Brief Description of the Drawings

Figure 1 is an amino acid sequence of a human intestinal trefoil factor (ITF; Accession No. BAA95531) (SEQ ID NO.:1).

Figure 2 is an amino acid sequence of a human pS2 protein (Accession No. NP_003216) (SEQ ID NO.:2).

Figure 3 is an amino acid sequence of human spasmodic polypeptide (SP; Accession No. 1909187A) (SEQ ID NO.:3).

Figure 4 is a cDNA sequence encoding a human intestinal trefoil factor (SEQ ID NO.:4).

Figure 5 is a cDNA sequence encoding a human pS2 protein (SEQ ID NO.:5).

Figure 6 is a cDNA sequence encoding a human spasmodic polypeptide (SEQ ID NO.:6).

Figure 7 is the nucleotide sequence of a gene encoding human intestinal trefoil factor (locus 10280533:52117-55412) (SEQ ID NO.:7).

Figure 8 is the nucleotide sequence of a gene encoding human pS2 protein (locus 10280533:16511-21132) (SEQ ID NO.:8).

Figure 9 is the nucleotide sequence of a gene encoding human spasmodic polypeptide (locus 10280533:957-5208) (SEQ ID NO.:9).

Detailed Description

5 The invention provides methods and compositions useful for the treatment, amelioration, and prevention of a wide range of lesions to the respiratory epithelium. Lesions of the respiratory epithelium treated according to the present invention can be caused by physical (e.g., surgical intervention or intubation), chemical (e.g., smoking or exposure to volatile solvent), or thermal trauma;
10 vascular compromise (e.g., resulting from congestive heart failure or chronic obstructive pulmonary disease); infective or inflammatory processes; antineoplastic therapy (e.g., radiotherapy or chemotherapy); or other diseases processes such as cystic fibrosis or asthma, for example. Furthermore, another common chemical insult to the respiratory epithelium includes the exposure to
15 high concentrations of oxygen (e.g., hyperbaric oxygen therapies) for extended periods of time.

Treatment of these lesions according to the invention can speed epithelial healing, reduce symptoms associated with the disruption to the airway epithelium, and reduce, delay or prevent the secondary complications of worsening rhinitis,
20 asthma, pneumonitis, or other complications of the airway epithelial lesion. Further, since the invention will speed normal epithelial closure and reduce infection, it will reduce the chance of both acquiring secondary infections as well as late secondary effects of ongoing sensitization of the airway (e.g., hay fever and asthma).

25 Lesions of the respiratory epithelium, such as those resulting from allergic reactions or from physical trauma, are amenable to trefoil peptide therapy delivered as an aerosol or a dry powder. The composition is formulated (micronized) into a dry powder inhaler, or an aerosol according to known and conventional methods for preparing such formulations. When used to treat the
30 tracheo-bronchial respiratory epithelium, administration of a composition of the

invention preferably occurs as soon as symptoms occur and will last on the order of three to ten days, or alternatively until the lesion to the respiratory epithelium disappears. In the case of milder lesions however, trefoil peptide therapy may resolve the lesion in a shorter period of time, particularly when combined with
5 another active ingredient.

The compositions of this invention can also be used prophylactically, prior to therapies that will damage the respiratory epithelium. For example, the compositions can be administered prior to anti-neoplastic therapy or prior to a surgical intervention in order to mitigate the loss of epithelial integrity.
10 Prevention or amelioration of symptoms due to nasal-pharyngeal respiratory epithelial disruption may also be achieved by administering the trefoil peptide prior to the anticipated insult. For example, a patient may be administered trefoil peptide therapy before the exposure to tree or grass pollen in "hay fever" season, or by administering prophylactic treatment at reduced intervals, during the period
15 when the patient is at risk for nasal-pharyngeal infections.

Typically, a metered dose inhaler or dry powder inhaler will be self-administered by the patient. Tidal breathing from a continuous nebulizer, usually under physician supervision, also allows for independent regulation of trefoil peptide and adjunct pharmaceutical dosages.

20

Pharmaceutical Formulations

Aerosols

Aerosolized formulations deliver high concentrations of the trefoil peptide directly to the airways with low systemic absorption, and include for example
25 nasal sprays, inhalation solutions, inhalation suspensions, and inhalation sprays. Nasal sprays typically contain a therapeutically active trefoil peptide dissolved or suspended in solution or in a mixture of excipients (e.g., preservatives, viscosity modifiers, emulsifiers, or buffering agents), in nonpressurized dispensers that deliver a metered dose of the spray. Inhalation solutions and suspensions are
30 aqueous-based formulations containing the trefoil peptide and, if necessary,

additional excipients. Such formulations are intended for delivery to the respiratory airways by inspiration. Typically, metered-dose aerosol inhalers create droplets that are 20 to 30 microns in diameter.

A major limitation of pulmonary delivery is the difficulty of reaching the deep lung. To achieve high concentrations of a trefoil peptide solution in both the upper and lower respiratory airways, the trefoil peptide is preferably nebulized in jet nebulizers, a ultrasonic nebulizer, or an electronic nebulizer particularly those modified with the addition of one-way flow valves, such as for example, the Pari LC PlusTM nebulizer, commercially available from Pari Respiratory Equipment, Inc., Richmond, Va., which delivers up to 20% more drug than other unmodified nebulizers.

The pH of the formulation is also important for aerosol delivery. When the aerosol is acidic or basic, it can cause bronchospasm and cough. The safe range of pH is relative and depends on a patient's tolerance. Some patients tolerate a mildly acidic aerosol, which in others will cause bronchospasm. Typically, an aerosol solution having a pH less than 4.5 induces bronchospasm. An aerosol solution having pH between 4.5 and 5.5 will occasionally cause this problem. The aerosol solution having a pH between 5.5 and 7.0 is usually considered safe. Any aerosol having pH greater than 7.0 is to be avoided as the body tissues are unable to buffer alkaline aerosols and result in irritation and bronchospasm. Therefore, the pH of the formulation is preferably maintained between 5.5 and 7.0, most preferably between 5.5 and 6.5 to permit generation of a trefoil peptide aerosol well tolerated by patients without any secondary undesirable side effects such as bronchospasm and cough. The osmolarity of the formulation can also be adjusted to osmolarities of about 250 to 350 mosm/L, according to the patient's tolerance. The administration of a hypertonic or a hypotonic solution may be poorly tolerated in certain instances, particularly when administered to a denuded mucosa. Propellants, such as HFA 134a, HFA 227, or combinations thereof, may also be used in the formulation. If desired, excipients that promote drug dispersion or enhance valve lubrication may also be formulated with the trefoil

peptide.

Dry powder formulation

As an alternative therapy to aerosol delivery, the trefoil peptide may also be administered in a dry powder formulation for efficacious delivery into the endobronchial space. Such formulations have several advantages, including product and formulation stability, high drug volume delivery per puff, and low susceptibility to microbial growth. Therefore, dry powder inhalation and metered dose inhalation are most practical when high amounts of trefoil peptide need to be delivered, including for example cases in which a large portion of the respiratory epithelium is affected with lesions. Depending on the efficiency of the dry powder delivery device, effective dry powder dosage levels typically fall in the range of about 20 to about 60 mg. The invention therefore provides a sufficiently potent formulation of a trefoil peptide in dry powder or metered dose form of drug particles. Such a formulation is convenient because it does not require any further handling such as diluting the dry powder. Furthermore, it utilizes devices that are sufficiently small, fully portable and tend to have a long shelf life.

For dry powder formulations of the invention, a trefoil peptide composition is milled to a powder having mass median aerodynamic diameters ranging from 1-10 microns by media milling, jet milling, spray drying, super-critical fluid energy, or particle precipitation techniques.

Particle size determinations may be made using a multi-stage Anderson cascade impactor or other suitable method. Alternatively, the dry powder formulation may be prepared by spray drying or solution precipitation techniques. Spray drying has the advantage of being the least prone to degrading the trefoil peptides. Solution precipitation is performed by adding a co-solvent that decreases the solubility of a drug to a uniform drug solution. When sufficient co-solvent is added the solubility of the drug falls to the point where solid drug particles are formed which can be collected by filtration or centrifugation. Precipitation has the advantage of being highly reproducible and can be performed under low temperature conditions, which reduce degradation. Super-

critical fluid technology can produce particles of pharmaceutical compounds with the controlled size, density and crystallinity ideal for powder formulations.

The dry powder formulations of the present invention may be used directly in metered dose or dry powder inhalers. Currently, metered dose inhaler
5 technology is optimized to deliver masses of 1 microgram to 5 mg of a therapeutic. Spacer technology, such as the aerochamber, may also be utilized to enhance pulmonary exposure and to assist patient coordination.

An alternate route of dry powder delivery is by dry powder inhalers. There are two major designs of dry powder inhalers, device-metering designs in which a
10 reservoir of drug is stored within the device and the patient 'loads' a dose of the device into the inhalation chamber, and the inspiratory flow of the patient accelerates the powder out of the device and into the oral cavity. Alternatively, dry powder inhalers may also employ an air source, a gas source, or electrostatics, in order to deliver the trefoil peptide. Current technology for dry powder inhalers
15 is such that payload limits are around 10 mg of powder. The dry powder formulations are temperature stable and have a physiologically acceptable pH of 4.0-7.5, preferably 6.5 to 7.0.

Therapeutic agents

20 In addition to the trefoil peptide, the therapeutic formulation according to the present invention may also comprise a second therapeutic agent, or regimen. The second therapeutic agent may be administered within (either before, or after administration of the trefoil peptide) 14 days, 7 days, 1 day, 12 hours, 1 hour, or simultaneously with the trefoil peptide. The second therapeutic agent can also be
25 present in the same or different pharmaceutical compositions as the trefoil peptide. Thus, pharmaceutical compositions for locally treating the respiratory epithelium may include, in addition to a trefoil peptide, for example, an anti-inflammatory compound, an antibiotic, a beta- adrenergic bronchodilator, a cholinergic receptor antagonist, a neurokinin receptor antagonist, a steroid, a
30 decongestant, a phosphodiesterase inhibitor, an analgesic, or an anesthetic.

When the second therapeutic agent is present in a different pharmaceutical composition, different routes of administration may be used. For example, the second therapeutic agent may be administered orally, or by intravenous, intramuscular, or subcutaneous injection. Thus, the second therapeutic agent need
5 not be administered by inhalation. If desired, more than one therapeutic agent may be administered with the trefoil peptide. Of course, pharmaceutical compositions may also contain two, three, or more trefoil peptides, or biologically active fragments.

10 *Trefoil Peptides*

The therapeutic trefoil peptide(s) are typically mammalian trefoil peptides or fragments thereof although non-naturally occurring homologs that are substantially identical to the mammalian trefoil peptides are also useful. Preferably, human trefoil peptides or fragments are used; however, trefoil
15 peptides from other species including rat, mouse, and non-human primate, may be used. Typically, the trefoil peptide is intestinal trefoil factor (ITF); however, spasmolytic polypeptide (SP), or pS2 are also useful.

Particular trefoil peptide fragments retain biological activity and may be substituted in any method or composition in which a trefoil peptide is used.
20 Methods and compositions containing a trefoil peptide, in which these trefoil peptide fragments may be substituted, are described, for example, in U.S. Patent Nos. 6,063,755 and 6,221,840, and U.S. Patent Application Nos. 10/131,363, filed April 24, 2002, 60/317,657, filed September 6, 2001, 60/327,673, filed October 5, 2002, 60/333,836, filed November 28, 2001, and 60/367,574, filed March 26,
25 2002 (hereby incorporated by reference).

Particularly useful ITF fragments that retain biological activity include the polypeptide corresponding to amino acid residues 15-73 of SEQ ID NO:1 (ITF₁₅₋₇₃) and amino acid residues 21-73 of SEQ ID NO:1 (ITF₂₁₋₇₃). Other useful ITF fragments are formed following cleavage of the C-terminal phenylalanine residue
30 (i.e., ITF₁₋₇₂, ITF₁₅₋₇₂, and ITF₂₁₋₇₂).

The biologically active trefoil peptide fragments of this invention can be produced using any appropriate method. For example, cDNA encoding the desired ITF fragment can be used with any method known in the art for producing recombinant proteins. Exemplary methods are provided herein. ITF fragments, particularly ITF₂₁₋₇₃, can be produced using a *Pichia* yeast expression system (see, for example, U.S. Patent Nos. 4,882,279 and 5,122,465) transformed with a cDNA encoding long ITF species, such as the full length ITF (e.g., SEQ ID NO: 4) or ITF₁₅₋₇₃, when the fermentation culture is maintained at pH ~ 5.0.

The trefoil peptides, including ITF, are soluble, and can therefore be dissolved in a pharmaceutically acceptable carrier liquid for aerosolization or nebulization for example. Aerosols containing a trefoil peptide are optimized for aerodynamic particle size, to target airway regions of interest. Typically aerosol sizes of 1-3 micron target deep lung (alveolar) structures, while a particle size of 5-10 micron result in tracheo-bronchial deposition. Moreover certain excipients may be used to prolong the local release of a trefoil peptide delivered in the lung or nasal region, or to retain the trefoil peptide formulation in the desired local area of the lung by modifying the mucociliary clearance rate.

Trefoil Peptide Dosages

Typically, the dosage, frequency and duration of therapy are tailored to the type and severity of the lesion being treated. For example, intermittent dosing may be sufficient to treat minor airway lesions. More severe airway lesions, resulting from, for example, severe smoke inhalation or thermal damage, may require continuous trefoil peptide administration. Alternatively, treatment may also be administered prophylactically, in anticipation of lesions to the respiratory epithelium. The prophylactic treatment may begin at least one day, three days, five days, seven days, or ten days prior to the insult. Treatment of unanticipated airway lesions preferably begin immediately after insult, or within 24 hours. Preferably, trefoil peptide therapy is administered at least one, two, three, four, or more than four times per day for at least one day, five days, fourteen days, or even

for the lifetime of the patient being treated. Alternatively, the trefoil peptide may be continuously administered to the patient over a set period of time, for a duration of one hour, two hours, 6 hours, one day, or more than one day for example. For this purpose, the trefoil peptide may be administered using a mask
5 adapter of a nebulizer system, for example.

Preferably, aerosol formulation contains a trefoil peptide concentration of 5, 10, 20, 40, 60, 80, 100 mg/mL, or more and is formulated in a physiologically acceptable solution, preferably in one quarter strength of normal saline. Ideally, the patient is administered with at least 10, 50, 100, 200, 500, 700, 1000, or more
10 than 1000 micrograms of a trefoil peptide administered as an aerosol. The use of dry powder inhalation preferably results in the delivery of at least about 1, 5, 10, 20, 30, 40, 50, 60, or more than 60 mg of the trefoil peptide to the respiratory airways of the patient receiving treatment. In such a formulation, the trefoil peptide is delivered as a powder in an amorphous or crystalline state in particle
15 sizes between 1 and 10 microns in mass median aerodynamic diameter necessary for efficacious delivery of the trefoil peptide into the endobronchial space for treatment, amelioration, and prevention of lesions of the respiratory epithelium. Fractions of 2 to 4 microns may also be employed to target the peripheral lung. Patient inspiration techniques, such as breath holding for example, may also
20 optimize deposition of the trefoil peptide.

If desired, the trefoil peptide may also be administered orally, or by intravenous injection, particularly in cases in which controlled or continuous release of the trefoil peptide is the goal.

All of the therapeutic agents employed in the compositions of the present
25 invention, including the trefoil peptide component, can be used in the dose ranges currently known and used for these agents. Different concentrations of either the trefoil peptide or the other agents may be employed depending on the clinical condition of the patient, the goal of the therapy (treatment or prophylaxis), the anticipated duration, the lesion site, and the severity of the damage for which the
30 trefoil peptide is being administered. Additional considerations in dose selection

include: disease etiology, patient age (pediatric, adult, geriatric), general health and comorbidity.

Anti-Inflammatory Agents

5 Any suitable anti-inflammatory agent can be formulated with the trefoil peptide and employed using the method of this invention. Suitable anti-inflammatory agents can be administered systemically, or can be administered by inhalation. Exemplary agents include, but are not limited to non-steroidal anti-inflammatory drugs (e.g., ibuprofen, tacrolimus, Cromolyn, Nedocromil),
10 cyclooxygenase-2-specific inhibitors such as rofecoxib (Vioxx®) and celecoxib (Celebrex®), and glucocorticoids.

Particularly effective glucocorticosteroid agents that may be used by aerosolization include for example beclomethasone, flunisolide, budesonide and triamcinolone. Other useful glucocorticosteroid agents include prednisolone,
15 dexamethasone and fluticasone. Although asthma is the main lung condition in which corticosteroids are used, such agents may also be useful when the respiratory epithelium is damaged by cigarette smoke as in chronic bronchitis and emphysema for example. Corticosteroids are also useful in the treatment of other lung diseases such as sarcoidosis, alveolitis and chronic inflammatory conditions.
20 These drugs may be given orally, intravenously (e.g., in severe cases), or by inhalation. Preferably, inhaled corticosteroids are administered to the patient because the dose required is much less and is delivered directly to the small air passages in the lungs with fewer associated side effects.

Anti-inflammatory concentrations known to be effective following
25 inhalation administration can be used. For example, ibuprofen may be present in the composition at concentrations sufficient to deliver between 25-800 mg per day to the respiratory lesion.

Bronchodilator Agents

Any active bronchodilator agent may be co-formulated with the trefoil peptide in the usual doses for respiratory application to the nasal-pharyngeal or tracheo-bronchial anatomy. Useful bronchodilators include, but are not limited to
5 methylxanthines (e.g., theophylline, theobromine, and caffeine), sympathomimetic agents (e.g., adrenaline, epinephrine, isoproterenol, and beta-adrenergic agonists), cholinergic receptor antagonists such as ipratropium bromide and tiotropium and neurokinin receptor antagonists.

Adrenergic bronchodilators are usually administered by inhalation to open
10 up the bronchial tubes (air passages) of the lungs and are typically used to treat, ameliorate, or prevent the symptoms of asthma, chronic bronchitis, emphysema, and other lung diseases. Such exemplary bronchodilators include albuterol, bitolterol, epinephrine, fenoterol, formoterol, isoetharine, isoproterenol, metaproterenol, pirbuterol, procaterol, racepinephrine, salmeterol, and terbutaline.

Alternatively, the trefoil peptide of the invention may be administered with
15 a leukotriene receptor antagonist (e.g., montelukast, or zafirlukast), a neurokinin receptor antagonist, an antihistamine (e.g., diphenhydramine, fexofenadine, cetirizine, or loratadine) or a cholinergic receptor antagonist.

Antimicrobial Agents

Any suitable antimicrobial agent can be used in the compositions of the invention at concentrations generally used for these agents. Suitable antimicrobial agents include, antibacterial, antifungal, antiparasitic, and antiviral agents. Exemplary antibacterial agents (antibiotics) include the penicillins (e.g., penicillin
25 G, ampicillin, methicillin, oxacillin, and amoxicillin), the cephalosporins (e.g., cefadroxil, ceforanid, cefotaxime, and ceftriaxone), the tetracyclines (e.g., doxycycline, minocycline, and tetracycline), the aminoglycosides (e.g., amikacin, gentamycin, kanamycin, neomycin, streptomycin, and tobramycin), the macrolides (e.g., azithromycin, clarithromycin, and erythromycin), the fluoroquinolones (e.g.,
30 ciprofloxacin, lomefloxacin, and norfloxacin), and other antibiotics including

chloramphenicol, clindamycin, cycloserine, isoniazid, rifampin, and vancomycin. Particularly useful formulations contain aminoglycosides, including for example amikacin, gentamicin, kanamycin, neomycin, netilmicin, paromomycin, streptomycin, and tobramycin.

5 Antiviral agents are substances capable of destroying or suppressing the replication of viruses. Examples of anti-viral agents include 1,-D-ribofuranosyl-1,2,4-triazole-3 carboxamide, 9->2-hydroxy-ethoxy methylguanine, adamantanamine, 5-iodo-2'-deoxyuridine, trifluorothymidine, interferon, adenine
10 arabinoside, protease inhibitors, thymidine kinase inhibitors, sugar or glycoprotein synthesis inhibitors, structural protein synthesis inhibitors, attachment and adsorption inhibitors, and nucleoside analogues such as acyclovir, penciclovir, valacyclovir, and ganciclovir.

 Antifungal agents include both fungicidal and fungistatic agents such as, for example, benzoic acid, undecylenic alkanolamide, ciclopirox olamine,
15 polyenes, imidazoles, allylamine, thicarbamates, amphotericin B, butylparaben, clindamycin, econazole, fluconazole, flucytosine, griseofulvin, nystatin, and ketoconazole.

 Other antimicrobial agents such as the antiparasitics like pentamidine, are known to have respiratory side effects. Therefore, co-administration of a trefoil
20 peptide and an antimicrobial of this type may reduce or prevent adverse events.

 Antimicrobial concentrations known to be effective in treating respiratory infections can be used.

Anticancer Agents

25 Cancers of the lung, including small cell and non-small cell carcinomas, damage the lung epithelium. Frequently, this injury is exacerbated by anticancer therapy because many anticancer agents have adverse effects on epithelial cells. Therefore, it is beneficial to administer trefoil peptide therapy in anticipation of, concurrent to, or following antineoplastic therapy to prevent, ameliorate, or treat
30 damage to the respiratory epithelium. Chemotherapeutics are usually

administered systemically by intravenous injection. The trefoil peptides may administered simultaneously, as an additive to the chemotherapeutic preparation, or separately, by inhalation. For patients undergoing radiation therapy, trefoil peptides are preferably administered by inhalation beginning one to three days
5 prior to each therapeutic session, continuing through the course of therapy, and continuing for one to three days after the final radiation treatment.

Production of Trefoil Peptides

Trefoil peptides and fragments can be produced by any method known in
10 the art for expression of recombinant proteins. Nucleic acids that encode trefoil peptides (e.g., human intestinal trefoil factor (Figure 4 and 7), human pS2 (Figure 5 and 8), and human spasmodic polypeptide (Figure 6 and 9) or fragments thereof may be introduced into various cell types or cell-free systems for expression thereby allowing large-scale production, purification, and patient
15 therapy.

Eukaryotic and prokaryotic trefoil peptide expression systems may be generated in which a trefoil peptide gene sequence is introduced into a plasmid or other vector, which is then used to transform living cells. Constructs in which the trefoil peptide cDNA contains the entire open reading frame inserted in the correct
20 orientation into an expression plasmid may be used for protein expression. Prokaryotic and eukaryotic expression systems allow for the expression and recovery of trefoil peptide fusion proteins in which the trefoil peptide is covalently linked to a tag molecule, which facilitates identification and/or purification. An enzymatic or chemical cleavage site can be engineered between the trefoil peptide and the tag
25 molecule so that the tag can be removed following purification.

Typical expression vectors contain promoters that direct the synthesis of large amounts of mRNA corresponding to the inserted trefoil peptide nucleic acid in the plasmid-bearing cells. They may also include a eukaryotic or prokaryotic origin of replication sequence allowing for their autonomous replication within the host
30 organism, sequences that encode genetic traits that allow vector-containing cells to

be selected for in the presence of otherwise toxic drugs, and sequences that increase the efficiency with which the synthesized mRNA is translated. Stable long-term vectors may be maintained as freely replicating entities by using regulatory elements of, for example, viruses (e.g., the OriP sequences from the Epstein Barr Virus genome). Cell lines may also be produced that have integrated the vector into the genomic DNA, and in this manner the gene product is produced on a continuous basis.

Expression of foreign sequences in bacteria, such as *Escherichia coli*, requires the insertion of a trefoil peptide nucleic acid sequence into a bacterial expression vector. Such plasmid vectors contain several elements required for the propagation of the plasmid in bacteria, and for expression of the DNA inserted into the plasmid. Propagation of only plasmid-bearing bacteria is achieved by introducing, into the plasmid, selectable marker-encoding sequences that allow plasmid-bearing bacteria to grow in the presence of otherwise toxic drugs. The plasmid also contains a transcriptional promoter capable of producing large amounts of mRNA from the cloned gene. Such promoters may be (but are not necessarily) inducible promoters that initiate transcription upon induction. The plasmid also preferably contains a polylinker to simplify insertion of the gene in the correct orientation within the vector. Mammalian cells can also be used to express a trefoil peptide. Stable or transient cell line clones can be made using trefoil peptide expression vectors to produce the trefoil peptides in a soluble (truncated and tagged) form. Appropriate cell lines include, for example, COS, HEK293T, CHO, or NIH cell lines.

Once the appropriate expression vectors are constructed, they are introduced into an appropriate host cell by transformation techniques, such as, but not limited to, calcium phosphate transfection, DEAE-dextran transfection, electroporation, microinjection, protoplast fusion, or liposome-mediated transfection. The host cells that are transfected with the vectors of this invention may include (but are not limited to) *E. coli* or other bacteria, yeast, fungi, insect cells (using, for example, baculoviral vectors for expression in SF9 insect cells),

or cells derived from mice, humans, or other animals. *In vitro* expression of trefoil peptides, fusions, or polypeptide fragments encoded by cloned DNA may also be used. Those skilled in the art of molecular biology will understand that a wide variety of expression systems and purification systems may be used to
5 produce recombinant trefoil peptides and fragments thereof. Some of these systems are described, for example, in Ausubel *et al.* (Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY 2000, hereby incorporated by reference).

Transgenic plants, plant cells and algae are also particularly useful for
10 generating recombinant trefoil peptides for use in the methods and compositions of the invention. For example, transgenic tobacco plants or cultured transgenic tobacco plant cells expressing a trefoil peptide can be created using techniques known in the art (see, for example, U.S. Patent Nos. 5,202,422 and 6,140,075). Transgenic algae expression systems can also be used to produce recombinant
15 trefoil peptides (see, for example, Chen *et al.*, Curr. Genet. 39:365-370, 2001).

Once a recombinant protein is expressed, it can be isolated from cell lysates using protein purification techniques such as affinity chromatography. Once isolated, the recombinant protein can, if desired, be purified further by e.g., high performance liquid chromatography (HPLC; e.g., see Fisher, Laboratory
20 Techniques In Biochemistry And Molecular Biology, Work and Burdon, Eds., Elsevier, 1980).

Polypeptides of the invention, particularly trefoil peptide fragments can also be produced by chemical synthesis using, for example, Merrifield solid phase synthesis, solution phase synthesis, or a combination of both (see, for example,
25 the methods described in Solid Phase Peptide Synthesis, 2nd ed., 1984, The Pierce Chemical Co., Rockford, IL). Optionally, peptide fragments are then be condensed by standard peptide assembly chemistry.

The following examples are intended to illustrate the principle of the present invention and circumstances when trefoil peptide therapy is indicated.
30 The following examples are not intended to be limiting.

Example 1: Treatment of Rhinitis due to Rhinovirus

The patient is administered a trefoil peptide-containing preparation beginning immediately after the onset of a head cold. The preparation contains a therapeutic dose of ITF₁₅₋₇₃. The trefoil peptide can be administered as a nasal spray using standard formulating methods to deliver 100 microliters of a 50 mg/ml spray of trefoil peptide. The patient receives medication by self-administering the nasal spray every 12 hours for the next five consecutive days. Also, the trefoil peptide active material may be applied with the standard dose of a nasal decongestant spray (e.g. 0.05% oxymetazoline HCl).

Example 2: Treatment of Allergic Rhinitis due to Grass Pollen

During hay fever season, the patient affected with allergic rhinitis is administered with antihistamines such as diphenhydramine, fexofenadine, cetirizine, or loratadine. Also, the patient is concurrently administered a nasal spray preparation containing a therapeutic dose of ITF₁₅₋₇₃. This component, in one example, is a nasal spray using standard formulating methods to deliver a 5 mg/ml spray of ITF. Continuing for the subsequent five days, the patient receives medication by self-administered nasal spray every 12 hours or as needed. In severe cases, the ITF active material may further be applied with the standard dose of a nasal glucocorticoid spray (e.g., beclomethasone, fluticasone, mometasone, or triamcinolone).

Example 3: Treatment of a Post Viral Prolonged Bronchospasm

In treatments for post-viral tracheo-bronchial epithelial disruption, the trefoil peptide containing material may be co-formulated with the standard dose of an inhaled salmeterol preparation, in a dry powder inhaler, an aerosol metered dose inhaler, or as a solution or a suspension in a ultrasonic or air-jet nebuliser. The treatment continues with the patient self-administering the medication every 12 hours for a period of at least 72 hours.

Example 4: Treatment of Adult Respiratory Distress Syndrome (ARDS)

Acute respiratory distress syndrome (ARDS) is a characteristic response of the lung in reaction to a wide variety of injury. Treatment of ARDS is initiated as soon as possible to minimize damage caused to the lung. The objective of

5 treatment is to provide enough support for the failing respiratory system (and other systems) until these systems have time to heal. The main supportive treatment of the failing respiratory system in ARDS is mechanical ventilation (a breathing machine) to deliver high doses of oxygen and a continuous level of pressure called PEEP (positive end-expiratory pressure) to the damaged lungs. To

10 speed healing, a trefoil peptide is administered by inhalation to patients with established ARDS or a syndrome of pre-ARDS. The amount of ITF₂₁₋₇₃ will be on the order of 1000 mg every 24 hours. The treatment is continued for at least 72 hours depending on the severity of the case and the clinical response of the patient. The regimen is repeated until healing or for ten days of therapy. It may

15 be more convenient to administer trefoil proteins to these patients less frequently (e.g. every 12 or 24 hours) and in higher concentrations with or without formulations to enhance the exposure of the lung capillary epithelium to the peptide. Additional forms of treatment that may be used along with the trefoil peptide therapy include for example antibiotics, immunosuppressants, blood

20 pressure supporting medications, tube feedings, and diuretics, which are used to reduce the fluid in the lungs. Since the pathology of ARDS is also linked to excessively produced nitric oxide, a NO blocker may be administered, if desired.

Example 5: Treatment of Human Respiratory Syncytial Virus

Human respiratory syncytial virus is the most important cause of

25 hospitalizations for viral respiratory tract disease in young children worldwide. Primary infection usually causes upper respiratory symptoms. Although the infection initiates in the upper respiratory tract, it can spread to the lower tract, via aspiration of secretions or via the respiratory epithelium, causing bronchiolitis and

pneumonia. During the infection, RSV causes extensive damage to the epithelium and the bronchiolar ciliary apparatus. Children affected by RSV may be administered ITF therapy to accelerate recovery of the respiratory epithelium. Patients are administered a trefoil peptide by inhalation, using for example, a dry powder inhaler, an aerosol metered dose inhaled, a solution or a suspension in a ultrasonic or air-jet nebuliser. The trefoil peptide is administered three times a day, at a dose of 1mg/puff. Desirably, Ribavirin, an aerosolized drug that can reduce the severity and the duration of illness, is also administered.

Example 6: Treatment of Influenza Infection

The influenza virus infects epithelial cells of the trachea and the bronchi. Extensive damage to the epithelium due to infection can cause severe coughing as well as pain in the chest, and the release of cytokines from damaged cells can further cause fever, chills, malaise, and muscular pains. Also, severe destruction of the mucous epithelium may lead to secondary bacterial infection and bronchitis. To alleviate the symptoms and accelerate the rate of recovery, the patient is administered trefoil peptide therapy as soon as symptoms of infection are manifested. ITF, or a biologically active fragment thereof, is administered in a dry powder inhaler, an aerosol metered dose inhaled, or as a solution or a suspension in an ultrasonic or air-jet nebuliser. Alternatively, patients may also be administered the trefoil peptide therapy by a nasal spray. This therapy is administered three to four times a day, and may be continued for a week following dissipation of the symptoms.

Example 7: Treatment of Chronic Bronchitis

Chronic Bronchitis is typically caused by chronic irritation of the respiratory airways or by microbial infections. As such, it is a condition often associated with smoking and its incidence is often associated with emphysema. Patients typically have a chronic cough with sputum. Damage to the epithelium from chronic bronchitis may predispose individuals to pneumococcal bacterial

invasion, which can lead to further complications, such as pneumonia. Therefore, restoration or improvement of the respiratory epithelium can alleviate symptoms associated with chronic bronchitis. Patients diagnosed with chronic bronchitis, or smokers, are immediately administered with a trefoil peptide in a dry powder
5 inhaler, an aerosol metered dose inhaled, or as a solution or a suspension in an ultrasonic or air-jet nebuliser. Patients can self-administer this regimen at least three times a day, for a period of at least seven days, or until the coughing ceases. If desired, the trefoil peptide therapy may also include administration of antibiotics.

10

Example 8: Treatment of Lesions Caused by Smoke Inhalation

Direct toxic effects caused by rapidly acting toxins such as smoke can incapacitate patients within moments. As such, the resulting effects, which include bronchospasm and alveolar damage, may cause rapid deterioration of the
15 patient and high mortalities. Inhalation of smoke can initiate an inflammatory response in a patient causing the release of histamine and other vasoactive substances that cause damage to the respiratory epithelium. Treatment will vary with the severity of the damage caused by smoke inhalation. The primary focus of treatment is to maintain an open airway and provide an adequate level of oxygen.
20 If the airway is open and stable, the patient may be given high-flow humidified 100% oxygen by mask. If swelling of the airway tissues is closing off the airway, the patient may require the insertion of an endotracheal tube to artificially maintain an open airway.

The patient is also immediately and continuously administered ITF₁₅₋₇₃ by
25 jet nebulizer for at least five days to reduce smoke-induced damage to the airway epithelium and the deleterious effects of hyperbaric oxygen therapies.

Example 9: Treatment of Asthma

The management of asthma is concerned primarily with the relief and prevention of symptoms through the treatment of underlying inflammatory processes, which cause damage to the respiratory epithelium. Furthermore, if
5 untreated, chronic inflammation makes the airways hyper-responsive to stimuli such as cold air, exercise, dust mites, pollutants in the air, thus exacerbating damage to the epithelium. Consequently, the asthmatic patient is administered with theophylline, an anti-inflammatory agent and a therapeutically effective amount of ITF₁₅₋₇₃ to ameliorate asthma-associated symptoms and to reduce
10 damage to the respiratory airways.

What is claimed is:

CLAIMS

1. A method for treating lesions of the respiratory epithelium in a mammal, comprising administering to said mammal a composition comprising a therapeutically effective amount of a trefoil peptide, or a biologically active fragment thereof.
2. The method of claim 1, wherein said trefoil peptide is intestinal trefoil factor (ITF), spasmolytic polypeptide, or pS2.
3. The method of claim 1, wherein said biologically active fragment is ITF₁₅₋₇₃, ITF₂₁₋₇₃, ITF₁₋₇₂, ITF₁₅₋₇₂, or ITF₂₁₋₇₂.
4. The method of claim 1, wherein said mammal is a human.
5. The method of claim 1, wherein said lesion is the result of an allergic reaction.
6. The method of claim 1, wherein said lesion is the result of asthma.
7. The method of claim 1, wherein said lesion is the result of a bacterial, viral, or fungal infection.
8. The method of claim 1, wherein said lesion is the result of inhalation of a chemical exposure, particulate matter, or smoke.
9. The method of claim 1, wherein said lesion is the result of a thermal burn.

10. The method of claim 1, wherein said lesion is the result of drug-induced lung damage, or anti-neoplastic therapy.

11. The method of claim 1, wherein said lesion is the result of trauma.

5

12. The method of claim 11, wherein said trauma is the result of a surgical procedure or intubation.

13. The method of claim 1, wherein said lesion is the result of chronic obstructive pulmonary disease or asthma.

10

14. The method of claim 1, wherein said lesion is the result of hyperbaric oxygen therapy.

15

15. The method of claim 1, wherein said administration is by inhalation.

16. The method of claim 15, wherein said composition is administered using a metered dose inhaler, or dry powder inhaler.

20

17. The method of claim 1, wherein said composition is an aerosol or a dry powder.

18. The method of claim 17, wherein said composition is nebulized in a jet nebulizer, ultrasonic nebulizer, or electronic nebulizer.

25

19. The method of claim 1, wherein said composition further comprises a second therapeutic agent.

20. The method of claim 19, wherein said trefoil peptide and said second therapeutic agent are administered in the same formulation.

30

21. The method of claim 19, wherein said trefoil peptide and said second therapeutic agent are administered by different routes of administration.

5 22. The method of claim 19, wherein said trefoil peptide and said second therapeutic agent are administered within 24 hours of each other.

23. The method of claim 19, wherein said second therapeutic agent is tobramycin.

10

24. The method of claim 19, wherein said second therapeutic agent is an anti-inflammatory agent, antimicrobial agent, antihistamine, neurokinin receptor antagonist, leukotriene receptor antagonist, decongestant, cholinergic receptor antagonist, phosphodiesterase inhibitor, or beta-adrenergic bronchodilator.

15

25. The method of claim 24, wherein said anti-inflammatory agent is a glucocorticoid.

26. The method of claim 25, wherein said glucocorticoid is
20 beclomethasone, flunisolide, budesonide, triamcinolone, prednisolone, dexamethasone, or fluticasone.

27. The method of claim 24, wherein said anti-inflammatory agent is a non-steroidal anti-inflammatory agent.

25

28. The method of claim 27, wherein said non-steroidal anti-inflammatory agent is ibuprofen, tacrolimus, cromolyn, nedocromil, refecoxib, or celecoxib.

30

29. The method of claim 24, wherein said beta-adrenergic receptor agonist is albuterol, bitolterol, epinephrine, fenoterol, formoterol, isoetharine, isoproterenol, metaproterenol, pirbuterol, procaterol, racepinephrine, salmeterol, or terbutaline.

5

30. The method of claim 24, wherein said antimicrobial agent is amikacin, gentamicin, kanamycin, neomycin, netilmicin, paromomycin, streptomycin, or tobramycin.

10

31. The method of claim 24, wherein said antihistamine is diphenhydramine, fexofenadine, cetirizine, or loratadine.

32. The method of claim 24, wherein said cholinergic receptor antagonist is ipratropium bromide or tiotropium bromide.

15

33. The method of claim 1, wherein said biologically active fragment of a trefoil peptide is encoded by an isolated nucleic acid molecule that hybridizes under high stringency conditions to a polynucleotide having the sequence of SEQ ID NO.: 4, SEQ ID NO.: 5, SEQ ID NO.: 6, SEQ ID NO.: 7, SEQ ID NO.: 8, or SEQ ID NO.: 9.

20

34. A pharmaceutical composition suitable for inhalation administration, wherein said composition comprises a trefoil peptide, or a biologically active fragment thereof, and a pharmaceutically acceptable carrier.

25

35. The composition of claim 34, wherein said trefoil peptide is intestinal trefoil factor (ITF), spasmolytic polypeptide, or pS2.

36. The composition of claim 34, wherein said biologically active
30 fragment is ITF₁₅₋₇₃, ITF₂₁₋₇₃, ITF₁₋₇₂, ITF₁₅₋₇₂, or ITF₂₁₋₇₂.

37. The composition of claim 34, wherein said composition is an aerosol or a dry powder.

5 38. The composition of claim 34, wherein said composition further comprises a second therapeutic agent.

39. The composition of claim 38, wherein said second therapeutic agent is tobramycin.

10

40. The composition of claim 38, wherein said second therapeutic agent is an anti-inflammatory agent, antimicrobial agent, antihistamine, cholinergic receptor antagonist, neurokinin receptor antagonist, leukotriene receptor antagonist, decongestant, phosphodiesterase inhibitor, or beta-adrenergic
15 receptor agonist.

41. The composition of claim 40, wherein said anti-inflammatory agent is a glucocorticoid.

20 42. The composition of claim 41, wherein said glucocorticoid is beclomethasone, flunisolide, budesonide, triamcinolone, prednisolone, dexamethasone, or fluticasone.

43. The composition of claim 40, wherein said anti-inflammatory agent
25 is a non-steroidal anti-inflammatory agent.

44. The composition of claim 43, said non-steroidal anti-inflammatory agent is ibuprofen, tacrolimus, cromolyn, nedocromil, refecoxib, or celecoxib.

30

45. The composition of claim 40, wherein said beta-adrenergic receptor agonist is albuterol, bitolterol, epinephrine, fenoterol, formoterol, isoetharine, isoproterenol, metaproterenol, pirbuterol, procaterol, racepinephrine, salmeterol, or terbutaline.

5

46. The composition of claim 40, wherein said antimicrobial agent is amikacin, gentamicin, kanamycin, neomycin, netilmicin, paromomycin, streptomycin, or tobramycin.

10

47. The method of claim 40, wherein said antihistamine is diphenhydramine, fexofenadine, cetirizine, or loratadine.

48. The method of claim 40, wherein said cholinergic receptor antagonist is ipratropium bromide, or tiotropium bromide.

15

49. A composition suitable for inhalation administration comprising a trefoil peptide or a biologically active fragment thereof, encoded by an isolated nucleic acid sequence that hybridizes under high stringency conditions with the sequence of SEQ ID NO.: 4, SEQ ID NO.: 5, SEQ ID NO.: 6, SEQ ID NO.: 7, SEQ ID NO.: 8, or SEQ ID NO.: 9.

20

FIGURE 1

1 MLGLVLALLS SSSAEYVGL SANQCAVPAK DRVDCGYPHV
41 TPKECNRGC CFDSRIPGVP WCFKPLQEAE CTF (SEQ ID NO.:1)

FIGURE 2

1 MATMENKVIC ALVLVSMLAL GTLAEQTET CTVAPRERQN
41 CGFPGVTPSQ CANKGCCFDD TVRGVPWCFY PNTIDVPPEE
81 ECEF (SEQ ID NO.:2)

3/12

FIGURE 3

1 EKPSPCQCSR LSPHNRTNCG FPGITSDQCF DNGCCFDSSV
41 TGVFWCFHPL PKQESDQCVN EVSDRRNCGY PGISPEECAS
81 RKCCFSNFIF EVPWCFFPNS VEDCHY (SEQ ID NO.:3)

FIGURE 4

```
1  atgctggggc tggtcctggc cttgctgtcc tccagctctg ctgaggagta cgtgggcctg
61  tctgcaaacc agtgtgccgt gccagccaag gacaggggtg actgcggcta ccccatgtc
121  accccaagg agtgcaacaa ccggggctgc tgctttgact ccaggatccc tggagtgcct
181  tgggtgtttca agccctgca ggaagcagaa tgcaccttct ga (SEQ ID NO.:4)
```


FIGURE 5

```
1   atggccacca tggagaacaa ggtgatctgc gccctgggcc tgggtgtccat gctggccctc
61  ggcaccctgg ccgaggccca gacagagacg tgtacagtgg cccccgtga aagacagaat
121 tgtgggttttc ctggtgtcac gccctcccag tgtgcaaata agggctgctg ttctgacgac
181 accgttcgtg gggccccctg gtgcttctat cctaatacca tcgacgtccc tccagaagag
241 gagtgtgaat ttttag (SEQ ID NO.:5)
```

FIGURE 6

```
1   atgggacggc gagacgcca gctcctggca gcgtcctcg tctggggct atgtgccctg
61  gcggggagtg agaaaccctc cccctgccag tgctccaggc tgagcccca taacaggacg
121 aactgcggct tccctggaat caccagtga cagtgttttg acaatggatg ctgtttcgac
181 tccagtgtca ctgggggtccc ctggtgtttc caccctcc caaagcaaga gtcggatcag
241 tgcgtcatgg aggtctcaga ccgaagaaac tgtggctacc cgggcatcag ccccgaggaa
301 tgcgcctctc ggaagtgctg cttctccaac ttcattttg aagtgccctg gtgtttcttc
361 ccgaagtctg tggaagactg ccattactaa
```

(SEQ ID NO.:6)

FIGURE 7

```
1   atgetggggc tggtcctggc cttgctgtcc tccagctctg ctgaggagta cgtgggcctg
61  tgtgagtact gccctgactg ccccggtggc aggggtgggc tgaagggag ggatccagga
121 taagggggga ttctgcattc atttaataat ggccacctgt cacatataca ctttttcctg
181 cgctagccct ttgaagtggg tctttattgt ccccatttca cagacaagga aaccgaggct
241 cagagaaagt taacaactta tccaaggcag ccctgcccag tctgtgttga aatcagggtt
301 tgagcctgag cccatcccct atgaccccat agccatcttt gctggagatt tctaaattac
361 aatataggtc tttatgcatt gttccacatt taaaaagaaa aaggaaagat gcaggagaaa
421 aaccctgact tcagaacact gtcaataccg gcaggcacia ggttcattta gccattgcat
481 agcaaccctg ccatggggtg tggctgtctc attaacccaa gtttgaagga atgagggcct
541 ggcttttatc tgggtgtctt ctgagcaggg tcaaaggcag tgggtcccga acttgagcc
601 cattagaatc acctggagag ctttaaaaaat cctaattgctt ggggcacacc agttacatca
661 gggcatctcc aggaagatc caggcctcag ctgttttgtt ttgagatagc cttgctttgt
721 cactcactgc tggagtgcag tggcacaatc tcagctcact gcaacctccg cctcctgggt
781 tcaagcaatt cttgtgcctc ggcttcaagt agctgggatt acaggcatgc accaccatgc
841 ccagctaatt ttttggattt ttagtagaga tggagtctcg ctatgttggc caagctggtc
901 tcaaactcct ggctcaagt gatcctcctg ccttggcctc ccaaagtgtc ggaattacag
961 gtgtaagcca ccatgcccag ccaacgtcag tcatttttaa agctctgcag ctgattccag
1021 tgtgagcgaa gtttggatgc caggaggata agcaattacg gactgggagc aagagaaggg
1081 aatgtaagac actgcacgtg attgccattt tcctaaggaa atactcagtt cgtaaatgaa
1141 acgcagtga cttctgtctg acatacagac atagaggctt gcctgaaaca tgaaaatatt
1201 ggggactgaa ggatgtcccg ggagggtggg acatgctcaa caattcagga aggggagatg
1261 cagaaaaaag tgaaaagcag gcagcatgcg ttgcaatgat ctctatggcg tgtgcctctc
1321 ctgtcacggg tttcatttaa aacaaaaggg caaggttttg ttggtcaaac aatgaagggt
1381 aactttgttt ctgggttcaa gggaccccag attcccaggg gggtcctgcc agctggaagg
1441 taccaggtc cgtagtgac ttcccagaaa ggtgataaga gcgtgccaa gagaagaca
1501 cttaggcaaa tggccagagt ccccgagctg agcatttaac agactgcctc tctttaaata
1561 ttcacaggga aagtgcattc tcctaagggc gagggtttca gcagtgggtg aactcggcgg
1621 ggtggggcgg agcgggagga tgcaaaactg caaagtgaag caaacacact caccgcagc
1681 cagcaagggc tctggcagct gacagggtt tgtctgggac agctgcaaac cagtgtgccg
1741 tgccagccaa ggacagggtg gactgcggct acccccatgt caccaccaag gagtgaaca
1801 accggggctg ctgctttgac tccaggatcc ctggagtgcc ttggtgtttc aagcccctgc
1861 aggaagcagg taaggcccca gtggcatcgt ggtctgggcc cagccccata aggcaggggg
1921 tctcagggcc tccctgtcct ttctgggctg gagatggagg cacaaggacc ccaggaagcc
1981 acacacacac acctgttcca aggcctcaga gcagaggctt cacacttagg gcagccatgg
2041 ccaggggctg tcctcttctg tccccttat gtaaaacata aaagcaattg tttcaaaaag
2101 gtgttcaaaa tgatggcatc gcatagaggg aactgattta gtaactattc ttgagagaag
2161 tggaaacgca taggtgtgga aagccgggcc gacttttggg ctgtttttgc aaatcgcccc
2221 cccagagtct tgcatttgtt ggcattcccct acacagacgg caggcgggtcc cagccctaga
2281 cgtaggcct cggtgccaca cccacctcc cccactctgc cccccacaag ggtcatctcc
2341 tctccctctc tctgccgtgg tggagggcag gtgcagggca accaccctgg gggttccctc
2401 cccaggggcg gagagcctgc gtgctgtgcg ggtaacagat ggccctgcac acgggtttgc
2461 caccctggct ccaccaggct tagctgcccc acatcgtggg tggggcgatt ggctataagc
2521 catctgcat gtccaagtgc cagctcagcc cccacgaagg ccgcacctgc gtgaggtacc
2581 ttcctggaac cagcatccag aggggcctct cttgcccttt gtccatagggt gaaatgcggg
2641 aggtgagtc ctgctggccc cggctccctg atcaatgatg ggcccctgcc cagggcctcc
```

FIGURE 7

```
2701 cttcaccctc cccagcaagt ccagggtagg ggtgggggtg ggggtccaga gaaggccagg
2761 agagagaggg gtctggctac tgtccactgc cggtcctgtt ccttcagctc cactggaact
2821 aactctcct ctgagtgcc gccatggccc tgccaaggcc catctcgctt gttatctgcc
2881 tgatccctgg gtcccactat cttgcttagc aaccgaggt gggaatcttg gctattcccc
2941 catgtggtgg ggactcaaca ctccccggtg actctgggga ggaggcagca ctagggtgctg
3001 gccttggagc ctgccctgac cttgggaagc tgggcagcgt gggaggagag agactgctca
3061 cacaagcctt tgctctgttt gcagaatgca cttctgagg cacctccagc tgcccccgcc
3121 cgggggatgc gaggctcgga gcacccttgc ccggctgtga ttgctgccag gcactgttca
3181 tctcagcttt tctgtccctt tgctccccgc aagcgcttct gctgaaagtt catatctgga
3241 gcctgatgtc ttaacgaata aagggtcccat gctccaccgc (SEQ ID NO.:7)
```

FIGURE 8

```
1 ccctggggtg cagctgagct agacatggga cggcgagacg cccagctcct ggcagcgctc
61 ctcgctcctg ggctatgtgc cctggcgggg agtgagaaac cctgtaagtg aaggagaggg
121 tctttttatg tgctttcttt atttctctta aagaaaaaaa aaaagcacia ccataaattt
181 acttgagagg ggggaatggct ataaaggcat ctggcaatgt gtgttggtca catgggattt
241 gccactgtctc aggagggtgg ctccaagaag ggcctccctc ctagggaag gctgagtgac
301 ggcaggtgtc agcgggcccc gtgtcgggcc aggaggcat tcccaccaag ggtccttgga
361 gtcccagagc actcacctct cgctggatc ttggccttg gtccatctgt tcacctcct
421 ctaggaggtt tttgtttttg tttttttccg agacaggatc tggctttgcc gcccaggcag
481 gagtgagtg gtgtgatctt ggctcactgc aacctctgcc tcccaggctc aagtgatcct
541 cccacctcag ccgcctgagt agctgaaacc acagtgtgtg accatcatgc ccggccaatt
601 tttttttttg tattgtttgt agagatgggg ttctgacatg ttgccagga ttgtcttgaa
661 tccctagcct ccagcaatct gccgcctcg gcttcctaaa gtgctgggat tataggtag
721 agccaccatg cctggctttt tttttttttt ccttttaaac taatataaca atttcagcaa
781 agccctatcg gcttctcagg aggaaaccgc attgcttaaa tatgggcaag ataagacttt
841 gtgtttctct atgtggcaac aagacagtag aggcattccc tagaacctct gagagaagga
901 gcagtgtggt ctgggggtacc aggggtggggc cgactgaggg tctttccaca gcccctgcc
961 agtgctccag gctgagcccc cataacagga cgaactgcgg cttccctgga atcaccagtg
1021 accagtgttt tgacaatgga tgctgtttcg actccagtgt cactgggggc ccctgggtgt
1081 tccacccccct cccaaagcaa ggtaatcttc cagggaatct tcttgggcca gctagtggca
1141 acccaggacc cagcttcaca ggcggagccc agagcagggg ccggaggagg cccagttgct
1201 agtctagggt tagcctgggt gggtagtct cgagctagcc ccggttggtt agtctggggc
1261 tagccaggt tggtagtct agagctagcc cagggttggtt agtctggggc tagccaggt
1321 tggtagtct ggggctagcc cagggttggtt agtctagggc tagttagggc tagttagtct
1381 aaggctagcc cagggttggtt agtttgagc tagcgcaggt tggtagtct ggggctagta
1441 gccaggttg gtagcctgg agctagccca ggttggttag tctagggcta gcgtaggctg
1501 gtagtctgg ggctagccca ggttggttag tctggagcta gccaggttg gctagtctgg
1561 ggctagtagc ccagggttggt tagtctgggg ctagcccagg ttggttagtc tagggctagt
1621 gtaggctagt tagtctaggg ctagcccagg ttagttagtt tggagctagc acagggtgat
1681 tagtctgggg ctagtagcct aggttggtta gtctggagct agcccaagtt ggtagtcta
1741 gggctagcat aggtctggtta gtctggggct agtagcctag gtttgtagt ctggagctag
1801 cccaggttg ttagtctagg gctagcgtag gctggttagt ctagggctag cccaggttg
1861 ttaatcggag ctagcccagg ttggttagtc tggagctagc ccaggttggt tagtctgagg
1921 ctagtagccc aggttggtta gtctggggct agcccaggtt tgtagtctg gagctagccc
1981 aggttggtta gtctggagct agcccaggtt ggtagtctg ggactagcct ggactgctag
2041 tctagaggta gcctagagga ctgctagtct agaggtagtc tagggctagc ccaggttggt
2101 tagtctgggg ctagcccag ttggttagtc ttagactagc ctggactgct agtctagagg
2161 tagccaggt tgtttagtct ggtactagcc tggactgcta gtctagaggt agcccaggt
2221 ggttaggttg gtagtctgg gactagctg gactgtagt ctagaggtag cccaggttg
2281 ttagtctggg actagcctgg actgtagtc tagaggtagc ccagattggt tagtctggga
2341 ctagtctgga ctgctagtct agaggtagcc caggttggtt agcctggggc cagcctggac
2401 tgtagtcta gaggtaacct aggtcagcca acagtgagat gaaaatttcc cactaccct
2461 gtttctacac tgttagttct ttcaacagac atgtgtgtgt ggagccatca gttttacttt
2521 agttgagaaa aaaatatata tatatatagt aggtctcttc tagtttttga agtgtgactt
2581 ctgaagaagc ttccatgggg aaatgaaggt atttaatagg acagcagtaa cataagggct
2641 gacagccctc aaatgttagg gaaggaagtg aagccttcta gggttctttg ggagttagtt
```

FIGURE 8

```
2701 ttatgttagt gcacgggac aggacccaag ttgtaacgcc gacgagtgct caaaggaagg
2761 ttgtgtgtgt gtcgtgcacc tgtgtgcgtg gaaccaggca cgtcctctgg agaaggagga
2821 ttcaccccca agattgttgc tgggaggcct gctgggcccc gcagggaaac caggcagatg
2881 gtggattgtt cactgaatgg cagtgtcttt gggaatcaat accatgtcca
2941 aacgctttcc atcttaccaa ggtgccca aaccttttct catcttggcc cgggggacca
3001 cccattttac tgagaacact gagtcccag aggcaaaatg atttcccaa ggcgggggac
3061 tccagagcct ctgactgtga ccaccccaca tgggcccac cttcgaggag gacaggccag
3121 ccaagcgtcg ctggggccga cacttccaca gtcccgggg gaggcgggcc caggggccga
3181 cacttccaca gtcccgggg gaggccgtcc cgggggatgc tgcccaggc agcacctcat
3241 gatccacgga ggctgcaa atcagcgtgct ctcagaggag gaagggtg agctttccag
3301 ggacacagcag gcctgactgg gtctcggtgc tgtgcctgtc ccatggcaga gtcggatcag
3361 tgcgtcatgg aggtctcaga ccgaagaac tgtggctacc cgggcatcag ccccgaggaa
3421 tgcgcctctc ggaagtgtcg cttctccaa ttcattcttg aagtgcctg gtgcttcttc
3481 ccgaagtctg tggaaagtaa cgtcgctgtg ggactctctg tctggttccc ggacaccatg
3541 attcctctc cgtccgtaga ggtgggggtg agggagggga gctgcctcgc agcctcagt
3601 ccatcgaggc caggggccct gcctcctat ggattctgaa ggcaattcca gaatgttctt
3661 ggcaaagaca gcgtcttttc aataagtta tagcctccag cattgccact gcgtcatctg
3721 tgatggctct agaaacagcg gctcatccct gttgcctccc cagggtgttg aacgttcaga
3781 ggcgttgctt gttttattgc aagcccatct gcatttgagg gctactgagt gtcttgact
3841 gtgctgggta ccagagagg ccactcaaa gcagacctgg ccccttctcc cgtggcttcc
3901 ccgttctccc ccacatgacc ccgaatgaca aacctcatcc acaacgtcct gctccgggca
3961 gtcccgggag ggtcccgccg gcagagggtga acgggtccac ttctcccacc cgttagtga
4021 tagtgtgttc ctgactcgga gtgtggcgag gtaaaaaaag accaagcaga tccaggaaaa
4081 tggggaaga gctactggcc cttgaaggat gccttttctt ttccttttgt taggatatca
4141 aagcactcca aagagcgaaa tatttcatgt tcaggatttt ccgagtgatt tttttatgt
4201 gacctaaagg tccacctaga aaatgttcac ttgtctgggg agaatgcgcc ccacagagga
4261 aactctggcc tgggggtggga agatttggtc cctttacacc cctccccgg gaaaggagct
4321 ccttcttcag taggaagctc ctgggcaaag tgatgcacgc ccacccagc ttcgcagcct
4381 aggcactccc atttctgggg ttcccttacc aaccatcttg catttaaact tctagactgc
4441 cattactaag agaggctggt tccagaggat gcatctggct caccgggtgt tccgaaacca
4501 aagaagaaac ttcgccttat cagcttcata cttcatgaaa tcctgggttt tcttaaccat
4561 cttttctca ttttcaatgg tttaacatat aatttcttta aataaaaccc ttaaaatctg
4621 ct (SEQ ID NO.:8)
```

11/12

FIGURE 9

```
1 atccctgact cggggtcgcc tttggagcag agaggaggca atggccacca tggagaacaa
61 ggtgatctgc gccctgggtcc tgggtgtccat gctggccctc ggcaccctgg ccgaggccca
121 gacaggtaag gcgtgcttct tcctgctctg tggggccaca gccagctctg gcagcctcog
181 ccaggagcca ctgtttttaca tacatatattt tgagcacctg ttttgtgcca ggtgctgttc
241 taggccctta aaagtatatc caattttacag gatcgccaaa agcagggtgga gagtaactca
301 ggggtggcagg gcccccgag accttcgaga agtgcgacga ggaggggggt gccttcagtc
361 ggggctgttt tcctgtgtta ggaagactat acaatcctcc caagtgtcat gtttcaaaga
421 ggaagtgttg gcgtggggtc tcagaatagt gcttttgact gttcatgcca acatctcccc
481 caggggcaga ccctcccaag gcccatccag ataggcccaa atgccgggtcc cagtgtggc
541 cacctgggag accctctccc acaggcccg atgcccgctc cagtgggtggc caactgggag
601 accctctcct acagggttctt gggctcccc gggatccatg ctctgggagt caaagccacc
661 tctctcatga gtgcgtggct ggcaacccat attccctggg gttgtcaagt ggatcggttg
721 ccctgggtcc ttctaggagg tgaggaggga ggccattctt gcttccttgg gaagtgtttg
781 catctcaact cctttacctg cagaatggat caacgggtctg ccctagggtc gtcaggaaat
841 gctgtgtggc agcatctgcg acttgcactt tgccagctgt ggggagctga ataacttatt
901 tgccgttatt aggtacagtt tcaagggtggg ggaggagaa agggctttct acgtttccaa
961 agcaagggtt tccagagagg cctgaagagg gagcgccag tgggtgctgtc cgtgccccca
1021 ctgccctcca gccacctctt gatctctgct gtgggggtacc gggcctgagg ggtgggcttg
1081 ggcagcgtga aagagcagcc agcattgggc tgcaagtggga agaccccaa gccatggca
1141 gggagcgggg gagctttgga acccgagaga ggaagtggcc tcggtgtaca gaacgaactg
1201 ggtgggtccc cgtgctggcc acccccaggc ccatctgctt gcgcccttgc cccacccca
1261 gccccagct ctgccccctg tgctgtggga tcacagaggc cgtggcaaac tccccctccc
1321 accccacaca ccctctggct caaggctcag agcgtctttg cgggtcactc aggtccatga
1381 tcctgttaca actgaaatct agaaaattgt gattacagtt tagtgattc gtgtgtggaa
1441 accattttcca tttattttcca tcatgcgaca aagacaaagc ggggtgggcaa gacagagtct
1501 gccggaggga gagcaccggg gctggaaatc ttcctccctg aggaggaaac ccccccgacc
1561 cccaggatga tgatccctcc tcaccacggg gcctctcttg acccccacag tgtcccgggg
1621 gtgggcgatg atcaccttca cgtcgcgatg gatccagacc ccaggagggc aaggttccca
1681 tggagctgc tgggcagcgg gagctgaaca cggatccttc ccagcaagcc aggaacactt
1741 tctccaaaga catctcgagg cagtccctga tagcaaagca gacaagagaa cagccccctc
1801 cggcctcccc tggggcgccc tcacctgagc cagtgtggcc agactgagtt cctccccctc
1861 tatgccccaa ggcagggaca gggaccggag ggtgctctgg gctcctcttt cccccctgc
1921 tgcaggctgt caaccaccag atcctaatag gttgctttct gagaccttg attccgcgga
1981 gctcagagcc tgaagctctg gtgttagaac ctcttgcata agatcctgcg gcagccccc
2041 gccagcccca tctgtccacg tgtcttcctc ctctagatcc ctttctctac tgcctgctt
2101 caagtgttt cacagcttgt accctctgtc ggctcctcct agaccacccc acccggtcct
2161 ctacaccttac ctgcaatggg tttccacctc ctgaacacac ctgggtctct ggaatggcct
2221 ttgcccctgc ggctccatct tcacctgggtg aacctcctcc tgcagggagc cccctgctt
2281 tgttcaacct gcttgtcatt ggctctctcg gggagtggcc taccctcggt gttaccctgg
2341 gcacctggg acgatggcct tgcgttgtct cgcacatgtt cttgccttct tcctccatca
2401 gatccttaga ctcttttttt tttttttttg agatggagtc ttgctctgtc actcaggctg
2461 gagtgcgaatg gtgcgatctt ggctcactac aacctctgcc tcctgggttc aagtgtattc
2521 cctgcctcag cctcccaagt agctgggatt acagacgtgt gccacaatgc ccgcctaatt
2581 ttttgtattt ttagtagaga tggggcttca ccattttggg caggctgggtc ttgaactcct
2641 gacctcaagt gattcacctc cttcagctc ccaaagtgtc gggattacag gcatgagcct
2701 gggcccagat atttagactc ttattaatga cttctctggt ttttaattct ggggtctctc
```

FIGURE 9

```
2761 cacctggcac agtgcctggc ttttgccatg ctagctccca cttctcatgc acacaaatgg
2821 tgctcagtaa atatttatgt attgagtaaa atttaataat catttggtga aattaaaaag
2881 tgaataaata agttacctag aaagatgcaa agtccacaaa cctggggcac cttgcatttt
2941 ccctgagcgt aatgtttgca catcaggatg tgaggaccac gtctccctct catgtcctga
3001 gggttttata tccgcctcac tggacagtgt ctgatgtcat tggagaagga agctggatgg
3061 gtgtgtgcat gataacatca aggaattcag cccacaactt actttgcttc ttacctgtgc
3121 actttcagag acgtgtacag tggcccccgc tgaaagacag aatttgtggtt ttccctggtgt
3181 cacgccctcc cagtgtgcaa ataagggtcg ctgtttcgac gacaccgttc gtggggtccc
3241 ctgggtgcttc tatcctaata ccacgcgacgt ccctccagaa ggtatggcct ttttatacga
3301 tgggttctga agatttagaa ttagttagaa aagtcattta agactacaga ggctctgac
3361 agcatcacca gctatgcctt tacacagagt cacggccgcc agtgggtggtg caatggggta
3421 gcctgagtcg ggctgcattc aggtccagga atagaaaggc agggctaagg gacttgggaa
3481 gaaacctgat tcccccccg cttctcttca catctctaac caaaagcctg ggaagagcca
3541 ctgttggtaa cgctttctag cttgcctagg atagaggggg aaggcatgac gaaatctgaa
3601 gacatttcat gtattctttt tttttttttt tttttgaaat ggagtctcgc tccgttgccc
3661 ctgagctgga gtgcaatggt gcgatcttgg ctactgcaa tctctgcctc ctgagttcaa
3721 cctcagcttc ctagtagctg agattacagg tgtgtgccac tacgcccagc taaatttttt
3781 ttgtattttt agtatagacg gggtttcacc atgttgcca gaccgtctt gaactcttga
3841 cctcaggtga tctgcccgcc tcagcctccc agagagctgg gattacaggc gtgagccacc
3901 gtgcccggct gacagttcat gttttctaaa gaatgtgcct atggatactt taaagtaaaa
3961 actctgtaat tgtttaaatg tgaaagaaaa tgtttatcct cactaaagca tctctttctc
4021 cctccccctc acccctgtag aggagtgtga attttagaca cttctgcagg gatctgcctg
4081 catcctgacg cgggtgccgtc cccagcacgg tgattagtc cagagctcgg ctgccacctc
4141 caccggacac ctcagacacg cttctgcagc tgtgcctcgg ctcacaacac agattgactg
4201 ctctgacttt gactactcaa aattggccta aaaattaaaa gagatcgata tt
```

(SEQ ID NO.:9)

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 June 2003 (05.06.2003)

PCT

(10) International Publication Number
WO 03/045332 A3

(51) International Patent Classification⁷: **A61K 39/00**,
38/03, 38/16

(21) International Application Number: PCT/US02/38258

(22) International Filing Date:
27 November 2002 (27.11.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/333,836 28 November 2001 (28.11.2001) US

(71) Applicant: **THE GENERAL HOSPITAL CORPORATION** [US/US]; 55 Fruit Street, Boston, MA 02114 (US).

(72) Inventor: **PODOLSKY, Daniel, K.**; 157 Edmunds Road, Wellesley, MA 02481 (US).

(74) Agents: **CLARK, Paul, T.** et al.; Clark & Elbing LLP, 101 Federal Street, Boston, MA 02110 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
24 July 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 03/045332 A3

(54) Title: METHODS AND COMPOSITIONS FOR TREATING LESIONS OF THE RESPIRATORY EPITHELIUM

(57) Abstract: This invention features methods of treating lesions of the airway epithelium by local or systemic administration of intestinal trefoil peptides. The intestinal trefoil peptide can be administered either alone or in combination with one or more therapeutic agents.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/38258

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 39/00, 38/03, 38/16
US CL : 424/185.1; 514/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
U.S. : 424/185.1; 514/12

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
MEDLINE, EMBASE, BIOSIS, SCISEARCH, CAPLUS on STN

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 6,063,755 A (PODOLSKY et al) 16 May 2000 (16.05.2000), column 3, lines 61-67.	1-2, 4-18, 33-35, 37, and 49
Y	US 5,783,416 A (THIM et al) 21 July 1998 (21.07.1998), column 7, lines 23-29, especially claim 14.	1-2, 4-18, 33-35, 37 and 49
Y	BABYATSKY, M.W. Oral Trefoil Peptides Protect Against Ethanol- and Indomethacin-induced Gastric Injury in Rats Gastroenterology February 1996, Vol 110. No. 2, pages 489-97, especially abstract.	1-2, 8, 10, 19-22, 24, 27-28, 34-35, 37, 40, and 43-44
Y	TRAN, C.P. Trefoil Peptide TFF2 (Spasmolytic Polypeptide) Potently Accelerates Healing and Reduces Inflammation in a Rat Model of Colitis Gut May 1999, Vol 44. No. 5, pages 636-42, especially abstract.	1-2, 8, and 34-35
Y	BEERS, M. H. in The Merk Manual of Diagnosis and Therapy Seventeenth Edition, 1999, pages 556-568, 1044-1050, and 1110-1113.	24-32, and 38-48

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

B earlier application or patent published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T

Later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

Z

document member of the same patent family

Date of the actual completion of the international search

10 May 2003 (10.05.2003)

Date of mailing of the international search report

27 MAY 2003

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US
Commissioner for Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450

Facsimile No. (703)305-3230

Authorized officer

Phuong Huynh

Telephone No. (703) 308-0196

Christina Chan

Form PCT/ISA/210 (second sheet) (July 1998)

DELLA MAE COLLINS
PARALEGAL SPECIALIST
TECHNOLOGY CENTER 1000